

658
INDEX

	PAGE
ALEXANDER, J. W., On the Class of a Covariant Tensor	245
ALEXANDER, J. W., and G. B. BRIGGS, On Types of Knotted Curves . .	562
ARWIN, A., Some Further Theorems Concerning the Formation of Chains	43
AYRES, W. L., Concerning Continuous Curves and Correspondences . .	396
AYRES, W. L., Note on a Theorem Concerning Continuous Curves . . .	501
BAILEY, H. W., The Summability of Single and Multiple Fourier Series	69
BLACK, H. L., A Cremona Group Isomorphic with the Group of the Twenty-seven Lines on a Cubic Surface	433
BRIGGS, G. B., and J. W. ALEXANDER, On Types of Knotted Curves . .	562
CAIRNS, W. D., Development of Functions in a System of Approximately Orthogonal Functions	503
CHURCH, A., On the Form of Differential Equations of a System of Paths	629
DARKOW, M. D., Determination of a Basis for the Integral Elements of Certain Generalized Quaternion Algebras	263
DAVIS, H. T., Derivation of the Fredholm Theory from a Differential Equation of Infinite Order	309
DICKSON, L. E., Ternary Quadratic Forms and Congruences	333
DINES, L. L., Note on Certain Associated Systems of Linear Equalities and Inequalities	41
DINES, L. L., On Positive Solutions of a System of Linear Equations .	386
DINES, L. L., On Completely Signed Sets of Functions	393
ETTlinger, H. J., On the Inversion of the Order of Integration of a Two Fold Iterated Integral	65
FITE, W. B., Periodic Solutions of Linear Differential Equations . . .	59
FLANDERS, D. A., Double Elliptic Geometry in Terms of Point, Order and Congruence	534
GARVER, R., Transformations of One Principal Equation into Another .	112
GARVER, R., Division Algebras of Order Sixteen	493
GEHMAN, H. M., Some Relations between a Continuous Curve and its Subsets	103
GLENN, O. E., A Theory of Integers in Relation to the Iteration of Algebraic Functions	368
GRAVES, L. M., On the Existence of the Absolute Minimum in Space Problems of the Calculus of Variations	153
GRONWALL, T. H., On the Existence and Properties of the Solutions of a Certain Differential Equation of the Second Order	355
HAZLETT, O. C., The Arithmetic of a General Algebra	92
INGRAHAM, M. H., Note on the Extensions of Groups to Obtain n -th Roots	53
JACKSON, D., A Problem in Minima	587
LANGFORD, C. H., Some Theorems on Deducibility	16
LANGFORD, C. H., Theorems on Deducibility. (Second Paper.)	459
LATIMER, C. G., On Certain Indefinite Quaternary Forms Representing All Integers	327
LEFSCHETZ, S., Correspondence between Algebraic Curves	342
McFARLAN, L. H., The Focal Point for the Problem of Lagrange with One Variable End Point	180
MARIA, A. J., Generalized Derivatives	419

	PAGE
MERRIMAN, G. M., Concerning the Summability of Double Series of a Certain Type	515
MICHAL, A. D., and T. Y. THOMAS, Differential Invariants of Affinely Connected Manifolds	196
MICHAL, A. D., and T. Y. THOMAS, Differential Invariants of Relative Quadratic Differential Forms	631
MILLER, N., On Related Maxima and Minima	117
OLSON, H. L., Linear Congruences in a General Arithmetic	237
PEPPER, E. D., Asymptotic Expression for the Probability of Trials Connected in a Chain	318
REYNOLDS, C. N., On the Problem of Coloring Maps in Four Colors, I.	1
REYNOLDS, C. N., On the Problem of Coloring Maps in Four Colors, II.	477
SHOOK, C. A., The Mixed Mean Function	171
SINGH, A. N., On Some Types of Non-differentiable Functions	472
STONE, M. H., Expansions in Bessel Functions	271
TAMARKIN, J. D., On Fredholm's Integral Equations Whose Kernels are Analytic in a Parameter	127
TAYLOR, J. H., Parallelism and Transversality in a Subspace of a General (Finsler) Space	620
THOMAS, J. M., Note on a Differential Equation	240
THOMAS, J. M., On Systems of Total Differential Equations	379
THOMAS, T. Y., The Replacement Theorem and Related Questions in the Projective Geometry of Paths	549
THOMAS, T. Y., and A. D. MICHAL, Differential Invariants of Affinely Connected Manifolds	196
THOMAS, T. Y., and A. D. MICHAL, Differential Invariants of Relative Quadratic Differential Forms	631
USPENSKY, J. V., On the Development of Arbitrary Functions in Series of Hermite's and Laguerre's Polynomials	593
VANDIVER, H. S., Application of Algebraic Number Theory to Congruences Involving Binomial Coefficients	330
VANDIVER, H. S., Transformations of the Kummer Criteria in Connection with Fermat's Last Theorem. (Second Paper.)	451
WEATHERBURN, C. E., On Lamé Families of Surfaces	301
WHYBURN, W. M., On the Green's Function for Systems of Differential Equations	291
WIENER, N., A New Definition of Almost Periodic Functions	365
WONG, B. C., Sextic Surfaces with a Double Septimic Curve	251

ERRATA

Vol. 27.

Page 245, line 9, for "Real Hermitian" read "Hermitian".

Page 248, line 3, for " $f^{(r-1)}(\lambda)$ must equal $\alpha(\lambda)$ " read " $\alpha(\lambda)$ is a factor of $f^{(i)}(\lambda)$ ($i = 0, 1, \dots, r-1$)".

Vol. 28.

Page 297, Omit all subscripts in (3.4), (3.5), (5.1).

Page 297, line 4 from foot, for "The i th component etc." read "This solution is $\int_a^b G \cdot \beta \, dt + u$ ".

Page 299, In (4.6), (4.7) interchange the subscripts of K .

Page 299, line 8 from foot, omit subscripts.

ON THE PROBLEM OF COLORING MAPS IN FOUR COLORS, I.*

BY C. N. REYNOLDS, JR.

Introduction. The problem of coloring in four colors the map of a simply connected closed surface has been reduced to the problem of coloring maps in which certain configurations, known as reducible configurations, are absent.† In this paper we shall develop some methods of so analysing the known geometric reductions of our problem as to discover and to prove some of their more important implications.

We shall use the reductions due to Kempe, Birkhoff and Franklin together with Errera's reduction of pairs of adjacent pentagons surrounded by hexagons. The reductions of Kempe, Birkhoff and Franklin are sufficient hypotheses for our final theorem that any irreducible map has at least twenty-eight regions. The use of one of Errera's reductions simplifies the proof of that theorem by eliminating much of the case making which would otherwise be necessary.

Our fundamental method will be a systematic study of a set of geometric operations which suffice to build any connected configuration of pentagons which may exist in an irreducible map. Under these operations certain numerical topological characteristics are found to undergo well defined increments. Linear relations between these increments imply homogeneous linear difference equations which yield certain homogeneous relations between our topological characteristics. These relations were originally found inductively from a study of particular configurations.

Secondly we shall prove synthetically certain inequalities between the topological characteristics of an irreducible map, making use of the linear relations mentioned above to reduce them to more serviceable forms.

We shall then apply these inequalities to the proof of the theorem that any irreducible map must have at least twenty-eight regions.

Finally we shall describe: (a) A map of twenty-eight regions which is irreducible with respect to the reductions of Kempe, Birkhoff and Franklin, but reducible with respect to the reduction due to Errera which we mentioned above. This map proves that what we may call the pre-Errera

* Presented, in part, to the American Mathematical Society, Feb. 28, 1925.

† cf. Franklin, *The four color problem*, Amer. Journ. of Math., 44 (1922), p. 42; also A. Errera, *Une contribution au problème des quatre couleurs*, Bull. Soc. Math. de France, 53, (1925), p. 42. Most of our references will be to these two papers.

4988-